

	Migraine	Tension type	Cluster	Other trigeminal autonomic cephalgias*	Cervicogenic	Medication overuse
<b>Frequency</b>	1-several/month	Variable	Every other day to 1–8 per day	Multiple daily	Daily for weeks	15 or more days in a month
<b>Severity</b>	Moderate–severe	Mild–moderate	Severe–very severe	Moderate–severe	Moderate–severe	Mild–severe
<b>Location</b>	Unilateral or bilateral	Bilateral	Unilateral	Unilateral in trigeminal distribution centred around eye	Posterior–unilateral	Bilateral (may mimic primary headache medication is used for)
<b>Nature of pain</b>	Throbbing/pulsating	Pressing/tightening	Severe unilateral pain	Severe – may be sometimes very brief	Ache radiating from neck/occipital region	Diffuse ache/pressure
<b>Visual aura</b>	+/- (present in a third of patients)	-	+/-	-	-	+/-
<b>Duration</b>	4–72 hours	Variable	15–180 minutes	Seconds – 30 min	Hours/constant	Hours/constant
<b>Nausea</b>	++	+/-	+/-	-	+/-	-
<b>Photophobia</b>	++	+/-	+ often ipsilateral	+/-	-	-
<b>Vomiting</b>	+	-	+/-	-	-	-
<b>Activity</b>	Keep still	Normal	Restless/pacing	Keep still or pace	Worse with neck movements/sleep & lifting	Normal
<b>Examination</b>	Normal	Tender points	Cranial autonomic features during attack	Cranial autonomic features during attack	Pain on neck movement and pressure	Normal
<b>Suggested treatment</b>	See following pages	See following pages	See following pages Refer all with new cluster	* Refer to neurology (routine)	See following pages	See following pages